

MthSc 440/640 Problem Set #4

Due 2/8/12

1. Consider the three vectors $v_1 = (1, -2, 3)$, $v_2 = (2, 1, 2)$, and $v_3 = (1, 2, -1)$. We want to find a *nonnegative* combination of these vectors (if one exists) that produces the vector $(1, 2, 1)$. For example using the weights $1/3, 1/6, 1/3$ produces the vector $(1, 1/6, 1)$, which however is not the required vector $(1, 2, 1)$.

(a) Formulate this problem as a Phase I LP. Clearly define your variables and explicitly show the corresponding matrices/vectors A , \mathbf{b} , \mathbf{c} and \mathbf{x} .

(b) Use the Phase I method to determine a solution, if one exists, or to demonstrate that there is no solution. At each iteration show the current \mathbf{x} , the objective function value, as well as the entering and leaving variables. For all calculations, solve the appropriate linear systems rather than compute inverses.

2. Consider the following LP problem:

$$\begin{aligned} \max z &= 16x_1 + 6x_2 + 7x_3 + 4x_4 \\ \text{s.t.} \quad &4x_1 + 2x_2 + 3x_3 - x_4 \leq 8 \\ &x_1 + 2x_2 + 5x_3 + 5x_4 \leq 11 \\ &3x_1 + x_2 + x_3 + 2x_4 \leq 5 \\ &x_1, x_2, x_3, x_4 \geq 0 \end{aligned}$$

(a) Convert this problem into standard equality form.

(b) Verify that the basic variables $\{x_1, x_3, x_6\}$ define an optimal basis. Determine the exact (fractional) coordinates for the corresponding solution \mathbf{x} as well as the optimal objective function value z .

(c) Determine the set of *all* optimal solutions (x_1, \dots, x_7) to this linear program. Express your answer in parametric form.